

[3,5].
See also [1,2,4].

References

- [1] Lorenz Halbeisen and Norbert Hungerbühler. ‘A geometric representation of integral solutions of $x^2 + xy + y^2 = m^2$ ’. In: *Quaestiones Mathematicae* (2019), pp. 1–15 (cit. on p. 1).
- [2] Axel Kohnert and Sascha Kurz. ‘Integral point sets over \mathbb{Z}_n^m ’. In: *Discrete Applied Mathematics* 157 (2006), pp. 2105–2117 (cit. on p. 1).
- [3] Donald E. Knuth. *Computers & Typesetting*. Vol. E: *Computer Modern Typefaces*. Reading, Mass.: Addison-Wesley, 1986 (cit. on p. 1).
- [4] Jozsef Solymosi and Frank De Zeeuw. ‘On a question of Erdős and Ulam’. In: *Discrete & Computational Geometry* 43.2 (2010), pp. 393–401. arXiv: 0806.3095_. URL: https://example.com/0%D1%80%D1%83%D1%81_ (cit. on p. 1).
- [5] Emma Sigfridsson and Ulf Ryde. ‘Comparison of methods for deriving atomic charges from the electrostatic potential and moments’. In: *Journal of Computational Chemistry* 19.4 (1998), pp. 377–395. DOI: [10.1002/\(SICI\)1096-987X\(199803\)19:4<377::AID-JCC1>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1096-987X(199803)19:4<377::AID-JCC1>3.0.CO;2-P) (cit. on p. 1).